



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

EXERCISES.

396

SHOW how to solve the simultaneous equations:

$$1) \quad x = y \sin(z + \alpha) = y \sin z - a = y \sin(z + \beta) - b;$$

$$2) \quad x(1 - \sin y) = a, \quad x[1 - \sin(y + \beta)] = b.$$

[*W. M. Thornton.*]

397

GIVEN two straight lines referred to rectangular coordinates; find geometrically an abscissa such that the sum of the squares of the two corresponding ordinates shall be a minimum.

[*R. A. Harris.*]

398

IF any curve, plane or gauche, be represented by means of the coordinates P , Q , which are so connected with a set of orthogonal and isothermal coordinates p , q that $P = \text{function } p$, $Q = \text{function } q$, then the angles made by this curve and the curves, $Q = \text{constant}$, $P = \text{constant}$, are

$$\tan^{-1} \frac{dq}{dp}, \quad \tan^{-1} \frac{dp}{dq},$$

respectively.

[*R. A. Harris.*]

399

IF y be any cubic function of x between the limits 0, h , its mean value can be expressed in an infinite number of ways by the formula

$$\frac{1}{6}(y_1 - 2y_2 + y_3)(1 + 2\sin^2\varphi) + \frac{1}{2}(y_3 - y_1)\sin\varphi + y_2,$$

where

$$x_2 = \frac{h}{\sqrt{2}} \sec\varphi \sin(45^\circ - \varphi), \quad x_3 = x_2 + \frac{1}{2}h \sec\varphi, \quad x_1 = x_2 - \frac{1}{2}h \sec\varphi.$$

[*W. H. Echols.*]

400

IF y be any cubic function of x , its mean value in any interval $X_2 - X_1 = L$ can be expressed in an infinite number of ways in terms of only two ordinates by the formula

$$\frac{3\sin^2\varphi}{1 + 2\sin^2\varphi} y_3 + \left[1 - \frac{3\sin^2\varphi}{1 + 2\sin^2\varphi} \right] y_2 + \frac{1}{18} L^3 \frac{1 - 4\sin^2\varphi}{4\sin^2\varphi} \tan\varphi,$$

where

$$x_2 = X_1 + \frac{1}{\sqrt{2}} L \sec\varphi \sin(45^\circ - \varphi), \quad x_3 = x_2 + \frac{1}{2} L \sec\varphi.$$

[*W. H. Echols.*]

401

DETERMINE the forms of surfaces which are such that the areas of parallel sections are cubic functions of their distances from a given section.

[*W. H. Echols.*]

402

TWO circular arcs are tangent to each other and to the sides a and b of a triangle ABC at the points A and B . Show that the difference of curvature of the arcs is least when their common tangent makes with a and b the angles $\frac{1}{2}(3B - A)$ and $\frac{1}{2}(3A - B)$, respectively.

[*W. H. Echols.*]

403

IF a triangle be inscribed in an ellipse, by drawing chords parallel to the tangents at the points α, β, γ ; prove that the sides are

$$c' = 2b' \sin(\gamma - \beta), \quad c'' = 2b'' \sin(\alpha - \gamma), \quad c''' = 2b''' \sin(\beta - \alpha),$$

where b', b'', b''' are the semi-diameters conjugate to those at the points α, β, γ . Also the area of the triangle $= 2ab \sin(\alpha - \gamma) \sin(\beta - \alpha) \sin(\gamma - \beta)$.

[*G. B. M. Zerr.*]

404

THE area of a rectangle circumscribing an ellipse is

$$\frac{4ab}{b'^2} [(a^2 + b^2) b'^2 - a^2 b^2]^{\frac{1}{2}},$$

where b' is the semi-diameter conjugate to that at the point of tangency of one of the sides.

[*G. B. M. Zerr.*]

405

A TRAPEZOID circumscribing an ellipse is formed by the tangents at the extremities, and the tangents parallel to a chord passing through the points whose eccentric angles are α and β . Prove that its area $= 2ab \tan \theta \operatorname{cosec}^2 \theta$, where $\theta = \frac{1}{2}(\alpha - \beta)$.

[*G. B. M. Zerr.*]

406

IF y be any cubic function of x the mean value of y in any interval L is*

$$y_m + \frac{L^2}{24d^2} (y_1 + y_2 - 2y_m),$$

wherein y_m is the ordinate at the middle of the interval and the ordinates y_1 and y_2 are at a distance d on either side of y_m .

If x_m be the abscissa of the middle point of the interval, then in the limit when d vanishes we also have for the mean ordinate of the cubic $y = f(x)$,

$$y_m + \frac{L^2}{24} f''(x_m). \quad [\text{W. H. Echols.}]$$

* A generalization of Newton's rule.